

Talking With a Partner About Erectile Difficulties

Why Honest Communication Can Be an Important Part of Erectile Dysfunction Care

By Dr. Nizamuddin Qasmi

Founder & Chief Physician, Saira Health Care

Focused Practice in Sexual Disorders & Infertility

BUMS – Hamdard University, Delhi

MD, CGO

Certificate in Infertility – MGBIMS, Delhi

Certificate in Urology – London, UK

Masters in Male Infertility – MasterHealthPro (HealthPro)

Integrated Sexual and Reproductive Health – ISRH, UNFPA

Introduction

One of the most difficult things for many men experiencing erectile difficulties is not the erection problem itself—it is deciding how to talk about it with their partner.

In my clinical practice, I regularly meet men who have been carrying this concern silently for weeks, months, and sometimes years. They may continue to love their partner, feel sexual desire, and want emotional and physical closeness, yet they gradually begin avoiding intimacy because they are afraid that an erection may not occur or may not last.

Unfortunately, silence can create another problem.

The partner may begin wondering:

“Is he no longer attracted to me?”

“Does he not love me anymore?”

“Is there someone else?”

“Have I done something wrong?”

At the same time, the man may be thinking:

“What if I fail again?”

“How will I explain this?”

“Will my partner think I am weak?”

“What if my marriage is affected?”

This is why I tell my patients that erectile difficulty should not automatically be treated as a private burden that a man must carry alone. When a couple has a supportive



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

relationship, honest and sensitive communication can become an important part of recovery.

Erectile dysfunction, commonly called **ED**, means a persistent or recurrent difficulty obtaining or maintaining an erection firm enough for satisfactory sexual activity. An occasional erection problem during periods of tiredness, anxiety, illness, emotional stress or excessive alcohol use does not automatically mean that a man has chronic erectile dysfunction. Persistent symptoms, however, deserve proper medical assessment.

Modern sexual-medicine guidelines increasingly emphasize that treatment should consider not only erection quality but also the patient's expectations, psychological well-being, relationship and sexual partner.

Erectile Difficulty Is Not Necessarily Loss of Love or Desire

This is perhaps the most important message that both partners need to understand.

An erection and sexual desire are related, but they are not the same thing.

A man may deeply love and desire his partner but still have difficulty achieving an erection.

An erection depends upon a complicated interaction between:

the brain

emotions

hormones

nerves

blood vessels

penile tissue

medications

physical health

sexual stimulation

relationship circumstances

Therefore, erectile difficulty should not immediately be interpreted as rejection, lack of attraction or loss of masculinity.



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

Many physical conditions can contribute to ED, including diabetes, hypertension, cardiovascular disease, obesity, hormonal disorders, neurological conditions, kidney disease and problems involving the male reproductive system. Certain medicines can also contribute. Anxiety, depression, stress, low confidence and negative body image may either cause erectile difficulty or make an existing physical problem worse.

This is one reason I advise couples:

Do not begin with blame. Begin with understanding.

Why Is Talking About Erectile Difficulties So Difficult?

Sexual health remains surrounded by embarrassment in many societies.

A man may have no difficulty telling a doctor that he has fever, diabetes, acidity or back pain, but mentioning erectile function can suddenly become extremely difficult.

There are several reasons.

Fear of Being Judged

Some men incorrectly associate erection quality with masculinity.

When an erection becomes unreliable, they may feel that something fundamental about their identity has changed.

This can produce shame, embarrassment and withdrawal.

But erectile dysfunction is a health condition—not a measure of someone's worth as a man.

Fear of Hurting the Partner

The man may worry that discussing ED will make his partner believe that attraction has disappeared.

Instead of explaining the problem, he begins avoiding sexual situations.

Unfortunately, unexplained avoidance may be more painful for the partner than an honest conversation.

Performance Anxiety

One unsuccessful sexual experience can sometimes become the beginning of a cycle.

The man remembers what happened.



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

Before the next sexual encounter he thinks:

“Will it happen again?”

His attention shifts away from pleasure and intimacy toward monitoring the erection.

Anxiety increases.

The erection becomes more difficult.

The next experience then reinforces the fear.

Over time, sexual activity may begin to feel like an examination rather than an intimate experience.

Stress and anxiety are recognized contributors to erectile difficulty, and psychological counseling or cognitive-behavioural approaches may be helpful when these factors are important.

Erectile Dysfunction Can Affect Both Partners

Although ED occurs physically in the male partner, its emotional effect can involve both members of the relationship.

Research has described erectile dysfunction as a **shared sexual concern of couples**, rather than something affecting only one person. Partner involvement may therefore be helpful during education, counseling and selection of treatment.

The partner may experience:

confusion

rejection

reduced confidence

fear that attraction has disappeared

frustration

anxiety about initiating intimacy

concern about the relationship

concern about fertility

worry about the man's health

The man experiencing ED may simultaneously develop:

performance anxiety



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

embarrassment

guilt

low confidence

avoidance of intimacy

frustration

sadness

fear of disappointing the partner

Without communication, both people may make incorrect assumptions about what the other person is thinking.

Does Communication Actually Matter?

Yes.

Communication by itself cannot correct every physical cause of erectile dysfunction. If a patient has severe diabetes-related vascular disease, testosterone deficiency, neurological damage or another medical condition, conversation alone is obviously not the complete treatment.

However, good communication can remove additional psychological and relationship pressure surrounding the problem.

A meta-analysis involving 48 studies found that better sexual communication was positively associated with several measures of sexual functioning, including erectile function and overall sexual function. The research shows an association rather than proving that conversation alone cures ED, but it supports what we often observe clinically: couples who can discuss sexual concerns openly may be better placed to manage them together.

How Should I Tell My Partner About My Erectile Difficulty?

There is no single perfect sentence.

What matters more is the attitude with which the conversation begins.

Choose the Right Time

I usually advise patients not to begin this discussion immediately after an unsuccessful attempt at intercourse.



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

At that moment both partners may already be emotional.

Choose a private, comfortable and neutral time when neither partner feels rushed.

You may say:

“I want to talk to you about something personal because our relationship is important to me.”

This immediately tells your partner that the purpose of the conversation is connection—not criticism.

Tell Your Partner What You Are Experiencing

Use simple language.

For example:

“Recently I have sometimes been having difficulty maintaining my erection. I still feel attracted to you, and this is not because I don't want intimacy with you.”

This can answer the question the partner may already be afraid to ask.

You do not need complicated medical terminology.

Be truthful.

Use “We” Instead of “Me Versus You”

Sexual difficulties become harder when each partner begins protecting themselves from the other.

Try to approach the condition as something that can be handled together.

Instead of:

“I have a problem.”

You may say:

“I would like us to understand what is happening and deal with it together.”

This small change in language can make an important psychological difference.

Reassure Your Partner About Attraction

When medically and emotionally true, say clearly:



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

“This does not mean that I have lost interest in you.”

Do not assume your partner automatically understands this.

Erectile function is often mistakenly interpreted as a direct indicator of attraction.

It is not.

Blood circulation, nerves, hormones, stress, medicines and many health conditions influence erections.

Allow Your Partner to Speak

The purpose is not simply to explain yourself.

Ask:

“How has this been making you feel?”

Then listen.

Your partner may have been afraid to raise the subject because they did not want to embarrass you.

They may also be worried about your health.

A calm conversation allows both people to replace assumptions with information.

What Should a Partner Avoid Saying?

Words can either reduce performance pressure or increase it.

Statements such as:

“Why can't you perform?”

“This never happened before.”

“What is wrong with you?”

“You are not interested in me anymore.”

“Prove that you still find me attractive.”

“A normal man should be able to do this.”

can increase shame and anxiety.

Sexual performance does not improve through humiliation.

A much better response is:



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

“We don't have to prove anything tonight. We can understand this together.”

That sentence can sometimes remove enormous pressure.

Stop Treating Every Sexual Encounter as an Erection Test

This is especially important when performance anxiety is involved.

If every moment of intimacy becomes focused on whether the penis is sufficiently firm, the mind remains in a state of evaluation rather than relaxation.

Couples can temporarily reduce pressure by broadening their understanding of intimacy.

Affection, touching, kissing, emotional connection and other mutually comfortable forms of intimacy can continue without making penetration the only definition of a successful sexual experience.

The objective is not to ignore ED.

The objective is to stop allowing ED to control every intimate moment.

The Performance-Anxiety Cycle

I often explain this cycle to patients:

One erection difficulty → worry about recurrence → increased monitoring during sex → anxiety → reduced arousal or erection → further loss of confidence → more anxiety during the next attempt.

When this pattern develops, treating only erection mechanics may not completely resolve the problem.

European Association of Urology guidance supports psychological approaches such as cognitive behavioural therapy when indicated, with partner involvement and medical treatment where appropriate.

This is why some patients benefit from an integrated approach rather than simply asking:

“Which tablet should I take?”

Erectile Difficulties Should Also Be Medically Evaluated

Communication with your partner is important, but please do not make the opposite mistake of assuming every erection problem is purely psychological.

Persistent erectile dysfunction can sometimes be the first visible sign of a broader health problem.

It may be associated with:

diabetes

hypertension

high cholesterol

obesity

cardiovascular disease

smoking

hormonal disorders

low testosterone in appropriate clinical circumstances

thyroid disorders

neurological problems

pelvic surgery or injury

kidney disease

Peyronie's disease

psychological stress

depression

anxiety

certain medications

excessive alcohol consumption

recreational drug use.

Erectile Dysfunction and Heart Health

This deserves special attention.

The blood vessels supplying the penis are small, and vascular disease that affects circulation can contribute to erectile dysfunction.



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

Current cardiovascular and sexual-medicine guidance considers vasculogenic ED an important cardiovascular risk marker. The Princeton IV Consensus and contemporary European guidance therefore emphasize cardiovascular risk assessment in appropriate men presenting with erectile dysfunction.

This does **not** mean that every man with ED has heart disease.

It means persistent unexplained erectile dysfunction should not simply be hidden or treated through self-medication.

Sometimes the sexual symptom gives us an opportunity to identify diabetes, hypertension, abnormal cholesterol or cardiovascular risk earlier.

What Happens During Medical Evaluation?

When a patient consults me for erectile difficulty, I believe the consultation should go much deeper than asking, “Can you get an erection—yes or no?”

A proper evaluation may include discussion of:

The erection problem

When did it begin?

Was the onset sudden or gradual?

Can you achieve an erection?

Does the erection disappear quickly?

Are morning or spontaneous erections present?

Does the problem happen every time or only occasionally?

Sexual health

Is sexual desire normal?

Is ejaculation normal?

Is orgasm normal?

Is there penile pain?

Is there penile curvature?

Is there premature ejaculation?

Are there fertility concerns?

General health



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

We may need to consider:

blood pressure

diabetes

weight

cholesterol

cardiovascular health

hormonal symptoms

smoking

alcohol consumption

sleep

physical activity

medications

Psychological and relationship factors

I may also ask about:

work stress

performance anxiety

depression

relationship tension

previous unsuccessful sexual experiences

fear of infertility

sexual confidence

Depending upon the history and clinical examination, laboratory investigations may include glucose assessment, lipid profile and testosterone evaluation, among other tests when clinically indicated. Contemporary European guidelines specifically recommend assessment of glucose, lipid profile and total testosterone in the basic evaluation of ED to identify potentially reversible or modifiable factors.

Treatment Should Be Based on the Cause

There is no single treatment that is ideal for every patient with erectile dysfunction.



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

The treatment plan depends upon:

the underlying cause

severity

duration

age

cardiovascular health

medications

hormonal status

psychological factors

relationship circumstances

fertility plans

patient preference

This is why buying an unidentified “sex power” medicine without examination can be a serious mistake.

Lifestyle Measures Are Part of Treatment

Improving general health can also support erectile health.

Depending on the individual, I may advise:

regular physical activity

weight management

smoking cessation

limiting excessive alcohol

improving sleep

controlling diabetes

controlling blood pressure

correcting abnormal cholesterol

nutritious eating

stress management

Lifestyle measures including physical activity, weight management and cardiovascular risk-factor control can improve erectile function in some men and are recommended as part of comprehensive ED management.

Medicines for Erectile Dysfunction

When appropriate, doctors may prescribe medicines known as **PDE5 inhibitors**, including medications such as sildenafil or tadalafil.

They improve the physiological erectile response by supporting blood flow to the penis during sexual stimulation.

However, they are not appropriate for everyone.

They should be used after consideration of medical history, cardiovascular status, other medicines and possible interactions.

Patients taking nitrate medicines for angina or certain cardiovascular conditions require particular attention because dangerous blood-pressure lowering can occur when nitrates are combined with PDE5 inhibitors.

Never stop or change prescribed heart, blood-pressure, psychiatric or other medication simply because you suspect it is contributing to erectile dysfunction. Discuss it with your treating doctor first.

Other established options for selected patients can include vacuum erection devices, injectable treatments and penile prosthesis surgery when conservative approaches are unsuccessful or unsuitable.

The Unani Perspective on Erectile Difficulties

As a physician trained in the Unani system of medicine with a focused clinical practice in sexual disorders and infertility, I believe it is important to explain the Unani understanding in language that patients can easily understand.

Classical Unani literature has discussed male sexual weakness under concepts such as **Zu'f-i-Bah**, while erectile weakness or penile flaccidity has also been described within this wider traditional framework.

Importantly, classical Unani physicians did not always view sexual weakness simply as a problem of the genital organ. Traditional descriptions considered general health, temperament, psychological influences, vital organs and reproductive function.

Standard Unani literature published through India's Central Council for Research in Unani Medicine describes psychological factors among the possible contributors to

sexual debility and includes addressing these psychological influences among the principles of management.

This aspect is particularly relevant to our present topic.

Even centuries ago, the broader concept recognized that the mind and sexual function can influence one another.

How I Use the Unani Approach in Erectile-Difficulty Patients

At Saira Health Care, my aim is not merely to give every man the same medicine.

The first step is to understand **why** erectile difficulty is occurring.

In suitable patients, an individualized Unani treatment approach may be used as part of a broader management programme intended to support general vitality, sexual health and the patient's underlying functional imbalance according to Unani principles.

Depending upon the individual case, I consider:

general health and physical strength

digestive health

sleeping pattern

emotional stress

libido

erection quality

ejaculation

semen and fertility concerns where relevant

associated chronic illnesses

diet

lifestyle

medication history

psychological and relationship factors

Treatment should be individualized rather than selected merely because a product is advertised as a “sexual medicine.”

What Does Scientific Research Say About Unani Treatment for ED?

This is an area where we should be both open-minded and scientifically responsible.

A randomized open-label clinical study published in 2025 evaluated oral and topical polyherbal Unani formulations in men with erectile dysfunction. Thirty-six participants were divided between two treatment groups, and improvement was reported in erection-related measures and International Index of Erectile Function scores.

The result is encouraging for further research, but the study was small, involved only 36 participants and had a relatively short treatment period.

Therefore, it would not be scientifically correct to use one small study to claim that all Unani medicines are proven cures for every form of erectile dysfunction.

Broader reviews of complementary and herbal therapies have also emphasized that the quality and quantity of evidence differ considerably between products and interventions.

My approach is therefore to respect traditional Unani knowledge while also respecting modern diagnostic science.

Traditional medicine and modern investigation do not need to be enemies.

A patient can receive an individualized Unani-oriented treatment plan while still having diabetes, cardiovascular risk, hormonal problems or other important medical causes properly investigated.

Never Take Unknown “Herbal Sex Medicines”

“Natural” does not automatically mean safe.

Patients frequently purchase unlabelled powders, capsules or so-called herbal sexual-enhancement products without knowing their ingredients.

Some products sold as natural sexual enhancers have been found to contain undisclosed pharmaceutical ingredients, and supplements may also interact with prescription medicines. Major medical sources therefore recommend discussing complementary medicines and supplements with a healthcare professional before use.

I strongly advise patients:

Do not hide your medicines from your doctor.

Tell your doctor about prescription medicines, herbal medicines, supplements and any product you are taking for sexual performance.



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

Saira Health Care's Approach to Erectile Difficulties

At **Saira Health Care**, sexual problems are approached with confidentiality, dignity and individualized clinical evaluation.

My focused practice is in **sexual disorders and infertility**, and I have spent years working with men and couples who often arrive frightened because they believe that an erection problem means their sexual life, marriage or fertility is permanently damaged.

Very often, the first thing these patients need is correct information.

Our approach is based on several principles.

1. Identify the Cause

We try to determine whether the dominant factors appear:

vascular

metabolic

hormonal

neurological

medication-related

psychological

lifestyle-related

relationship-related

or, as happens frequently, a combination of several factors.

2. Treat the Patient, Not Just the Symptom

I do not believe that every patient who reports ED should automatically receive the same treatment.

Age, general health, sexual history, relationship circumstances, fertility goals and accompanying sexual symptoms all matter.

3. Give Attention to Psychological Factors

If a man can achieve normal erections in certain situations but repeatedly loses his erection when intercourse is attempted, performance anxiety deserves careful consideration.



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

This does not mean that the problem is “imaginary.”

Psychological processes produce genuine physiological changes.

The fear itself can interfere with sexual response.

4. Involve the Partner When Appropriate

If both partners are comfortable, partner involvement can sometimes make treatment easier.

The partner can understand:

what ED actually means

why it occurs

how treatment works

what improvement realistically looks like

how unnecessary pressure can be reduced

Modern ED guidance similarly encourages discussing the needs and expectations of both the patient and sexual partner.

5. Use Unani Treatment Responsibly

Where an Unani treatment approach is considered appropriate, it should be selected after proper consultation and individualized assessment.

Unani care should not delay investigation of important underlying disorders such as uncontrolled diabetes, serious cardiovascular disease, significant hormone abnormalities or neurological disease.

This combined clinical thinking is especially important in sexual medicine.

What Can a Partner Do to Help?

If your partner tells you that he is experiencing erection difficulties, remember that telling you may already have required significant courage.

You can help by:

listening without ridicule

avoiding blame

reassuring him that intimacy is not a performance examination

encouraging proper medical evaluation

reducing pressure surrounding intercourse

participating in counseling when appropriate

discussing sexual preferences openly

focusing on affection and connection

avoiding comparisons with previous experiences or other people

You do not need to become your partner's doctor.

You simply need to remain a partner.

What Should the Man Do?

If you are the person experiencing erectile difficulties:

Do not withdraw without explanation.

Your partner may misunderstand silence.

Do not repeatedly “test” yourself.

Constantly checking erection quality can increase anxiety.

Do not self-medicate indefinitely.

Persistent ED deserves evaluation.

Do not assume the condition is permanent.

Many causes of erectile dysfunction are treatable or manageable.

Do not blame yourself.

A medical problem is not a moral failure.

Take responsibility for seeking appropriate care.

Reassurance from a partner is helpful, but proper diagnosis remains important.

A Simple Conversation You Can Use

If you do not know how to begin, you can say:



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

“I want to tell you something because you are important to me. Recently I have sometimes been having difficulty getting or maintaining my erection. It is worrying me, but I don't want you to think it means I am not attracted to you. I would like us to approach it calmly, and I am planning to speak with a doctor so that I can understand the reason.”

Notice what this statement does.

It:

identifies the issue

reassures the partner

removes blame

shows responsibility

communicates affection

introduces a practical next step

That is often much more helpful than months of silence.

When Should You Consult a Doctor?

Seek professional evaluation when:

erectile difficulty occurs repeatedly

the problem continues for several weeks or months

erections have progressively become weaker

morning erections have significantly changed

you have diabetes

you have hypertension or cardiovascular disease

sexual desire has fallen significantly

you suspect low testosterone or another hormonal problem

you have penile pain or curvature

you have infertility concerns

you are experiencing severe anxiety or depression

erectile difficulty began after starting a medicine



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

the condition is affecting your relationship

you repeatedly rely on sexual-enhancement medicines without medical assessment

Persistent ED may be an opportunity to identify another medical condition rather than something that should simply be hidden.

When Is Urgent Medical Attention Required?

Seek urgent medical care if an erection remains painfully persistent for approximately four hours or longer.

This may represent **priapism**, which requires prompt medical treatment to reduce the risk of permanent tissue injury.

Sudden serious symptoms such as chest pain, severe breathlessness, fainting or other cardiovascular symptoms associated with sexual activity also require appropriate urgent medical evaluation.

Erectile Difficulty and Fertility Are Not the Same Thing

Another common misunderstanding I see is that erectile dysfunction automatically means infertility.

It does not.

Erectile function and sperm production are different biological processes.

A man can have excellent semen parameters but erectile difficulty.

Another man can have normal erections but severe abnormalities in sperm count, motility or morphology.

However, severe ED can indirectly interfere with natural conception if vaginal intercourse and semen deposition cannot occur reliably.

When a couple is also experiencing infertility, both sexual function and reproductive health should therefore be assessed separately.

This distinction is particularly important in my practice because I work with both **sexual disorders and male infertility**.

Can Erectile Difficulties Be Treated Successfully?

In many patients, yes, significant improvement is possible.



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

But I prefer the word **treatment** rather than giving every patient a promise of “permanent cure.”

The outcome depends upon the cause.

A young man with predominantly performance anxiety requires a different treatment approach from a man with longstanding diabetes and vascular disease.

A patient with medication-related ED requires a different strategy from someone with testosterone deficiency.

A patient with relationship stress requires something different from a man with severe arterial insufficiency.

Good medicine begins with this distinction.

Frequently Asked Questions

Can stress really cause erectile difficulty?

Yes.

Stress and anxiety can interfere with sexual arousal and erections, and existing ED can itself generate additional anxiety.

If I still have morning erections, does that mean there cannot be a physical problem?

No.

Morning or spontaneous erections provide useful clinical information, but they do not by themselves prove that the cause is purely psychological.

Your complete medical and sexual history should be assessed.

Should I tell my wife or partner?

In a supportive relationship, I generally encourage open communication.

Partner involvement is also supported by contemporary sexual-medicine guidance when appropriate.

Should I immediately take sildenafil?



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

Not without appropriate medical advice.

The correct treatment depends upon the cause, your medical history and medicines you are already taking.

Can diabetes cause ED?

Yes.

Long-term high blood glucose can damage blood vessels and nerves important for normal erectile function. Diabetes is an important risk factor for ED.

Does erectile dysfunction mean my partner no longer satisfies me?

No.

ED cannot be interpreted simply from attraction.

Physical and psychological health both contribute to erections.

Can couples counseling help?

Yes, particularly when performance anxiety, relationship tension, communication problems or psychological distress are contributing factors.

Counseling may be combined with medical treatment.

Can Unani medicine be used for erectile difficulties?

Unani medicine has a longstanding traditional approach to male sexual weakness, and emerging clinical research is evaluating specific Unani formulations.

In my practice, Unani care is individualized after assessing the patient's health and possible causes of ED.

However, responsible Unani treatment should not replace necessary investigation for diabetes, cardiovascular disease, hormonal abnormalities or other significant medical causes. Current Unani-specific clinical evidence remains promising but limited and requires larger, high-quality studies.

My Message to Patients



+91-9452580944



+91-5248-359480



www.sairahealthcare.com

If you are reading this because you recently experienced difficulty getting or maintaining an erection, please remember one thing:

Do not allow one difficult sexual experience to define your entire sexual health.

The human sexual response is not a machine.

Sleep, stress, emotions, health, hormones, blood circulation, medication, relationship circumstances and psychological expectations can all influence it.

If the problem repeats, do not hide it for months out of embarrassment.

Talk calmly with your partner.

Seek professional assessment.

Do not purchase unknown medicines merely because somebody promises instant sexual power.

And most importantly, do not turn intimacy into an examination that you have to pass every time.

At Saira Health Care, my objective while treating sexual disorders is not simply to improve a laboratory report or produce a temporary erection. My aim is to understand the individual patient as completely as possible—his physical health, reproductive health, psychological state, sexual confidence, lifestyle and relationship circumstances—and then provide a rational, individualized treatment strategy.

Modern medicine gives us powerful diagnostic tools and established treatments.

The Unani system gives us a longstanding tradition of individualized assessment, lifestyle regulation and therapeutic approaches to sexual health.

Used responsibly, with proper diagnosis and realistic expectations, patient-centered care can bring these perspectives together while always keeping safety first.

And sometimes, one of the first steps toward recovery is surprisingly simple:

Talk to your partner.

Not with shame.

Not with blame.

But with honesty.

Because erectile difficulty should not automatically become a wall between two people. With understanding, proper medical evaluation and appropriate treatment, it can instead become a health concern that both partners learn to face together.



+91-9452580944



+91-5248-359480



www.sairahealthcare.com



About the Author

Dr. Nizamuddin Qasmi

Founder & Chief Physician – Saira Health Care
Focused Practice in Sexual Disorders & Infertility

Dr. Nizamuddin Qasmi has focused his clinical work on the evaluation and management of male sexual disorders, reproductive-health concerns and infertility, with particular emphasis on individualized Unani treatment supported by appropriate clinical evaluation.

Professional Qualifications & Training

BUMS – Hamdard University, Delhi

MD

CGO

Certificate in Infertility – MGBIMS, Delhi

Certificate in Urology – London, UK

Masters in Male Infertility – MasterHealthPro (HealthPro)

Integrated Sexual and Reproductive Health – ISRH, UNFPA

Through **Saira Health Care**, Dr. Qasmi works to improve awareness around sexual disorders and infertility, reduce the stigma that prevents patients from seeking medical advice, and provide individualized care for patients who frequently suffer in silence because of embarrassment surrounding sexual-health conditions.

Important Medical Disclaimer

This article is intended for patient education and general health awareness. It is not a substitute for individual diagnosis, examination or medical treatment. Erectile dysfunction can have cardiovascular, metabolic, hormonal, neurological, psychological, medication-related and other causes. Patients with persistent symptoms should consult a qualified healthcare professional.

Do not start, stop or change prescription medication or sexual-enhancement products solely on the basis of online information.

Individual response to Unani, conventional or other treatments varies, and no treatment should be presented as a guaranteed cure for every patient.



+91-9452580944



+91-5248-359480



www.sairahealthcare.com